

Condensation of a lattice gas in three dimensions

Johannes Zierenberg*, Micha Wiedenmann, Wolfhard Janke

Institute for Theoretical Physics, Universität Leipzig, Leipzig, Germany

*zierenberg@itp.uni-leipzig.de

We investigate the condensation transition of a lattice gas model [1, 2, 3]. This simplified model is restricted to nearest-neighbor interaction. For fixed density it is equivalent to the Ising model at fixed magnetization. In order to investigate the temperature dependence, we apply a novel parallel multicanonical method [4]. By this means, the diffusion properties of the particles can be adjusted in such a way that free-energy barriers can be circumvented. Keeping the density fixed, the condensation transition is studied as a function of system size. Our results may further be compared to an analytic prediction of equilibrium droplet formation [1]. To this end, we choose a fixed temperature and compare the average largest droplet to a rescaled density. Above a critical density, a single droplet is observed that includes more than half of the particle excess, consistent with theory.

References

- [1] M. Biskup, L. Chayes, R. Kotecký: *On the formation/dissolution of equilibrium droplets*. Europhys. Lett. **60**, 21–27 (2002)
- [2] A. Nußbaumer, E. Bittner, T. Neuhaus, W. Janke: *Monte Carlo study of the evaporation/condensation transition of Ising droplets*. Europhys. Lett. **75**, 716–722 (2006)
- [3] A. Nußbaumer, E. Bittner, W. Janke: *Monte Carlo study of the droplet formation-dissolution transition on different two-dimensional lattices*. Phys. Rev. E **77**, 041109 (2008)
- [4] J. Zierenberg, M. Marenz, W. Janke: *Scaling properties of a parallel implementation of the multicanonical algorithm*. Comput. Phys. Comm. **184**, 1155–1160 (2013)